

Urinary excretion of *in vivo* ¹³C-labelled milk oligosaccharides in breastfed infants

Silvia Rudloff^{1,2}, Gottfried Pohlentz³, Christian Borsch², Michael J. Lentze⁴ and Clemens Kunz^{2*}

¹Department of Pediatrics, Justus-Liebig-University Giessen, Feulgenstrasse 12, D-35392 Giessen, Germany

²Institute of Nutritional Science, Justus-Liebig-University Giessen, Wilhelmstrasse 20, D-35392 Giessen, Germany

³Institute for Medical Physics and Biophysics, University of Münster, Robert-Koch-Strasse 31, D-48149 Münster, Germany

⁴Department of Pediatrics, University of Bonn, Adenauerallee 119, D-53113 Bonn, Germany

(Submitted 12 April 2011 – Final revision received 10 June 2011 – Accepted 24 June 2011 – First published online 5 September 2011)

Abstract

Recent observations indicate that human milk oligosaccharides (HMO) are involved in a variety of physiological processes in infants. Their metabolic fate, however, is virtually unknown. We investigated metabolic aspects in infants after endogenous ¹³C-labelling of HMO. An oral bolus of natural and ¹³C-labelled galactose (Gal; 23 g Gal + 4 g ¹³C-Gal) was given to ten lactating women. Aliquots of milk at each nursing as well as breath samples from the mothers and urine from their infants were collected over 36 h. The ¹³C-enrichment of HMO and their renal excretion was determined by isotope ratio-MS; characterisation was achieved by fast atom bombardment-MS. After the Gal bolus was given, an immediate ¹³C-enrichment in milk and in infants' urine was observed which lasted 36 h. Mass spectrometric analysis of ¹³C-enriched urinary fractions confirmed the excretion of a variety of neutral and acidic HMO without metabolic modification of their structures. Components with glucose split off at the reducing end were also detectable. Quantitative data regarding the infants' intake of lacto-*N*-tetraose and its monofucosylated derivative lacto-*N*-fucopentaose II ranged from 50 to 160 mg with each suckling, respectively; renal excretion of both components varied between 1 and 3 mg/d. Since the intake of individual HMO by the infants was in the range of several hundred mg per suckling, i.e. several g/d, and some of these components were excreted in mg amounts as intact HMO with the infants' urine, not only local but also systemic effects might be expected.

Key words: Milk oligosaccharides: *In vivo* ¹³C-labelling: Infants: Urinary excretion

Human milk contains a large variety of complex oligosaccharides in concentrations ranging from 10 to 20 g/l^(1,2). The quantity of these components does not only depend on the lactational stage of the mother but is also affected by the expression of specific glycosyltransferases in the mammary gland^(1–3). Genes encoding for H and Lewis a or x antigens as well as the secretor status determine the presence of α1,2-, α1,3- and/or α1,4-fucosylated core structures of oligosaccharides. In addition, different patterns of sialylation, i.e. the attachment of α2,3- and/or α2,6-linked *N*-acetylneuraminic acid (NeuAc), increase the variability of human milk oligosaccharide (HMO) to a number of about 115 structures characterised so far^(4,5). The biological significance of HMO for the infant, however, has not been proven yet. Since the 1950s, HMO have been thought to be growth-promoting factors for the so-called 'bifidus flora' in the gut of breastfed infants⁽⁶⁾. Recently, the analysis of the genome of

Bifidobacteria indicated their evolutionary adaptation to preferentially use specific milk components, particularly HMO as their substrate^(7,8). However, the bifidogenic effect of HMO themselves and their direct impact on the intestinal microbiota are difficult to demonstrate *in vivo*. The same applies to other specific *in vitro* functions of HMO, such as their potential to influence inflammatory and infectious processes via inhibition of the attachment of pathogens to epithelial cells or leucocyte endothelial and neutrophil platelet interactions^(9–12). Animal and preclinical studies indicate that oligosaccharides like those in human milk may play an important role in many physiological processes such as influencing cell recognition and cell signalling, cell adhesion, the composition of the microbiota or even affecting neurodevelopment^(13–15).

To better understand the underlying mechanisms, more information regarding the metabolic fate of HMO in the infants

Abbreviations: FAB-MS, fast atom bombardment-MS; Gal, galactose; HMO, human milk oligosaccharides; HPAEC-PAD, high pH anion exchange chromatography with pulsed amperometric detection; IR-MS, isotope ratio-MS; LNFP II, lacto-*N*-fucopentaose II; LNT, lacto-*N*-tetraose; NeuAc, *N*-acetylneuraminic acid; NeuAcLac, *N*-acetylneuraminyl-lactose; PDB, Pee Dee Belemnite.

* **Corresponding author:** Professor C. Kunz, fax +49 641 9939049, email clemens.kunz@ernaehrung.uni-giessen.de

is needed. In a previous study⁽¹⁶⁾, we demonstrated that the application of stable isotopes to lactating mothers leads to the preferential incorporation of ¹³C-galactose (Gal) into lactose and milk oligosaccharides. The ¹³C-enrichment of HMO in this pilot study was high enough to investigate the metabolic fate of HMO in the infants. In an earlier study, we showed in one mother and her infant that the combination of fast atom bombardment-MS (FAB-MS) and isotope ratio-MS (IR-MS) is suitable to address specific questions with regard to the metabolism of HMO in humans⁽¹⁷⁾. In the present study, we report the renal excretion of intact and partly degraded complex oligosaccharides in ten infants after *in vivo* labelling of HMO by administering ¹³C-Gal to their mothers.

Subjects and methods

Subjects and study design

Exclusively breastfeeding women (*n* 10; 3–6 months post-partum) participated in the present study. Three mothers had already participated in our previous study⁽¹⁶⁾ and seven were newly recruited. Some dietary restrictions, i.e. the avoidance of naturally ¹³C-rich foodstuffs (e.g. corn, pineapple, fish), were given. The mothers were asked not to drink any milk for breakfast on the study day and to record their daily food intake during the study period. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the ethics committee of the University of Giessen. Written informed consent was obtained from all subjects/patients.

Between 08.00 and 09.30 hours, immediately after breakfast, mothers orally ingested a Gal bolus consisting of 23 g Gal + 4 g ¹³C-Gal dissolved in about 50 ml of drinking water. The purity of ¹³C-Gal (D-Gal; 1-¹³C) from Eurisotop (Saint-Aubin Cedex, France) was determined to be higher than 99%.

Milk and breath sampling of the women

A milk sample (5–10 ml) was collected immediately before the Gal bolus was given (baseline value) and at the beginning of each nursing during the following 36 h⁽¹⁶⁾. At the same time, mothers took breath samples and collected their urine in 3–4 h fractions.

Sampling of the infants' urine

For urinary collections, diapers consisting purely of cellulose and free of other absorptive material (Procter & Gamble, Frankfurt, Germany) were used. Urine was collected in adhesive bags which were emptied or replaced before each nursing. Diapers were changed before each nursing and immediately frozen at –20°C. For urine extraction, diapers were thawed and mechanically pressed leading to urine yields of about 70–80%. In previous experiments, it was verified that the urine collection via diapers and adhesive urine bags did not affect HMO analysis.

Separation of milk fractions and characterisation of milk carbohydrates

The separation of whole milk into fractions with fat, proteins and carbohydrates followed by the removal of lactose was achieved by ultracentrifugation, acetone treatment and gel filtration chromatography of the carbohydrate fraction as described earlier⁽¹⁸⁾. Then, neutral and acidic HMO were characterised by chromatography and MS (see the following subsections).

High pH anion exchange chromatography with pulsed amperometric detection

High pH anion exchange chromatography with pulsed amperometric detection (HPAEC-PAD) on a Carbo Pac PA-1 column (250 × 4.6 mm inner diameter) equipped with a guard column and a Model PAD 2 detector (Dionex, Sunnyvale, CA, USA) were used for characterisation of neutral and acidic HMO at the following conditions: eluent A, 100 mM-NaOH; eluent B, 100 mM-NaOH and 250 mM-sodium acetate. The elution programme started with 3 ml of buffer A followed by a gradient of up to 100% buffer B in 30 min and a re-equilibration volume of 5 ml buffer A. The flow-rate of 1.0 ml/min was used and 25 µl of 1 or 4 mg/ml solutions were injected. Molar response factors were determined by injecting three to six times equimolar amounts of each oligosaccharide as described previously⁽¹⁸⁾.

Fast atom bombardment-MS

Oligosaccharides (50–100 µg) were dissolved in 500 µl of pyridine–acetic anhydride (1:1, v/v). The mixture was stirred at room temperature overnight. Then the solvent was evaporated under N₂ and the residue was used for mass spectrometric analysis without further purification. The FAB-MS analysis was carried out on a VG-ZAB-T four-sector instrument (Fisons Instruments, Manchester, UK); caesium was used for atom bombardment⁽¹⁶⁾. The applied acceleration voltage was 8 kV. Thioglycerol–*m*-nitrobenzylalcohol (1:1, v/v) was used as matrix to which 1 µl of a homogeneous sample solution in chloroform–methanol (1:1, v/v) was added. For spectra recording, positive ions (FAB+) were detected on the second photomultiplier point detector. The spectra were run in a mass range from 100 to 3000 atom mass units with a scan rate of 8 s per decay and up to ten scans were accumulated.

Determination of ¹³C enrichment by isotope ratio-MS

¹³C enrichment was determined as δ¹³C_{PDB} measured in duplicate in whole milk, milk carbohydrates, total urine and urinary carbohydrate fractions by IR-MS (Isoprime; Isoprime Limited, formerly: GV Instruments; Manchester, UK) after total combustion at 1020°C (EuroEA, Eurovector, Milan, Italy)⁽¹⁶⁾. Breath samples were analysed in a BreathMat IRMS Microgas IRMS, Isoprime Limited). ¹³C elimination rates were then quantified as described in the following subsection.

¹³C analysis by isotope ratio-MS

The ¹³C content of a sample is expressed as delta value ($\delta^{13}\text{C}_{\text{PDB}}$) (‰) which is the relative difference between the ¹³C:¹²C ratio of a sample (Sa) and the international Pee Dee Belemnite (PDB) standard with a (¹³C:¹²C)_{PDB} ratio of 0.0112372,

$$\delta^{13}\text{C}_{\text{PDB}} (\text{‰}) = ({}^{13}R_{\text{Sa}} - {}^{13}R_{\text{PDB}}) / ({}^{13}R_{\text{PDB}}) \times 1000$$

where ${}^{13}R_{\text{Sa}} = ({}^{13}\text{C}/{}^{12}\text{C})_{\text{Sa}}$ is the isotope ratio of the unknown sample and PDB is Pee Dee Belemnite.

Data evaluation was performed with MassLynx inorganic version 4 software (GV Instruments) taking into account the ¹⁷O-content of the sample and the blank correction for the carbon of the tin cups⁽¹⁶⁾. For an easier comparison of the data, the results were expressed as $\Delta\delta^{13}\text{C}_{\text{PDB}}$ (corrected for the ¹³C:¹²C ratio at baseline). The accepted standard deviation was <0.2‰ for total urine and isolated carbohydrates and <0.7‰ for total milk. The accepted deviation for breath sample analyses was <0.3‰.

Calculation of the cumulative ¹³C-enrichment in milk and urine

To determine the ¹³C-enrichment in milk and urine, total organic carbon (TOC) and ¹³C-content (at%) for each sample were measured. Then, the excreted amounts of ¹³C were calculated, taking into account the baseline ¹³C values and the ¹³C-bolus given.

Using this ¹³C-at%_{sa} and the TOC of a sample, the absolute amount of ¹³C of each sample (Sa) was then calculated as follows:

$${}^{13}\text{C}_{\text{sa}} (\text{g}) = \text{TOC}_{\text{sa}} (\text{g}) \times \frac{{}^{13}\text{C-at}\%_{\text{sa}}}{100},$$

where at% is atom%.

Calculation of the cumulative ¹³C exhalation

The percentage recovery of the ingested ¹³C amount in breath (¹³C_{cum}) was calculated using the equation shown in the previous subsection. Thereby, the exhaled amount of ¹³C is defined as

$${}^{13}\text{C}_{\text{excreted}} (\text{mol/kg}) = \Delta\delta^{13}\text{C}_{\text{cum}} (\text{‰} \times \text{h}) \times {}^{13}R_{\text{PDB}} \times \text{CO}_2 - P (\text{mmol/kg} \times \text{h}),$$

with $\Delta\delta^{13}\text{C}_{\text{cum}}$ being the cumulative ¹³C exhalation.

Determination of galactose and lactose content

Gal and lactose contents of milk and urine were photometrically determined using a colorimetric kit from Boehringer Mannheim (Mannheim, Germany).

Calculation of ingested milk volumes and respective amounts of human milk oligosaccharide

To determine the amount of milk ingested by the infants, the milk volume was determined by weighing the infants on a digital electronic balance before and after each nursing.

Oligosaccharides were quantified by HPAEC-PAD analysis. Peak areas of lacto-*N*-tetraose (LNT) and lacto-*N*-fucopentaose II (LNFP II) were then calculated as percentage of the total amount of oligosaccharides applied, corrected by the molar response of the amperometric detector⁽¹⁸⁾.

Results

¹³C-enrichment of whole milk and ¹³CO₂ exhalation

In all milk samples, there was an immediate increase in ¹³C-enrichment of whole milk in the first hours after the oral ¹³C-Gal bolus was given to the mothers followed by a second smaller ¹³C-peak in six out of ten milk samples from the next morning (Fig. 1). To get an indication of how much Gal was metabolised for energy production, ¹³C-enrichment of breath was determined by analysing ¹³CO₂. Exhalation over 36 h compared to the ¹³C:¹²C ratio in whole milk resulted in a similar course although with a much higher enrichment in most samples (Fig. 1).

HPAEC-PAD analysis of the neutral and acidic fractions after Sephadex G25 gel filtration chromatography revealed a rather complex mixture of carbohydrates which were then characterised by FAB-MS. As an example, the composition of the major acidic and neutral HMO is given in Table 1. The molecular weight of these oligosaccharides varied between about 500 and 2500 Da.

Determination of the ¹³C-enrichment of the infants' urine

The ¹³C:¹²C ratio of urine fractions collected over 36 h after the Gal bolus was given to their mothers is shown in Fig. 1. The ¹³C-enrichment follows about the same pattern as in whole milk. Also, the first ¹³C-appearance in most urine fractions is delayed as compared to the corresponding milk fraction.

To ensure that the higher ¹³C:¹²C ratio is not due to the excretion of free ¹³C-Gal, we determined the total Gal excretion by enzymatic methods. The data which are exemplified for one infant in Fig. 2 revealed that the excretion of free Gal was only high in the first sample after the Gal bolus was given to the mother, and returned to baseline afterwards. In contrast to the total Gal excretion, the ¹³C-enrichments in the same urine fractions reached their maximum value 10–11 h (fraction 4) after the Gal bolus was given to the mother.

To characterise ¹³C-enriched carbohydrates, urine fractions were subjected to Sephadex gel filtration to obtain components of different molecular sizes and to separate monosaccharides and lactose from oligosaccharides. These fractions were further characterised by HPAEC-PAD, IR-MS and FAB-MS. The major components in these fractions are shown in Table 2. Not only small oligosaccharides such as the lactose derivatives Fu₁Lac and Fu₂Lac were detected but also LNT, one of the main core structures in HMO (Table 2). Moreover, the mono-, di- and trifucosylated derivatives of LNT, lacto-*N*-hexaose, lacto-*N*-octaoses and lacto-*N*-decaose could be found. Also, besides neutral components, *N*-acetylneuraminyl-lactose (NeuAcLac), the major acidic component in HMO and the sialylated derivative of LNT,




Fig. 1. ^{13}C -enrichment of whole milk ($-\diamond-$) and infants' urine ($-\diamond-$) and $^{13}\text{CO}_2$ exhalation by the mothers ($-\triangle-$) during the first 36 h after the oral intake of a galactose (Gal) bolus consisting of 4 g ^{13}C -Gal + 23 g Gal by the mothers. The $\delta^{13}\text{C}_{\text{PDB}}$ (‰) values of each sample are corrected over the baseline values of each respective sample obtained at time point 0, immediately before the Gal intake. The bolus was taken after breakfast which varied between 08.00 and 09.30 hours.

Table 1. Fast atom bombardment-MS of the major neutral and acidic human milk oligosaccharides from one mother in fractions after Sephadex 25 chromatography

Nominal m/z values of detected molecular ions			
$[M + H]^+$	$[M + Na]^+$	Oligosaccharide	Composition
Neutral peracetylated oligosaccharides			
	701	Lac	Hex ₂
	931	FucLac	FucHex ₂
1254	1276	LNT	Hex ₃ HexNAC
1484	1506	Fuc ₁ LNT	FucHex ₃ HexNAC
1714	–	Fuc ₂ LNT	Fuc ₂ Hex ₃ HexNAC
1829	–	LNH	Hex ₄ HexNAC ₂
2059	2081	Fuc ₁ LNH	FucHex ₄ HexNAC ₂
2289	2311	Fuc ₂ LNH	Fuc ₂ Hex ₄ HexNAC ₂
2634	2656	Fuc ₁ LNO	FucHex ₅ HexNAC ₃
2864	2886	Fuc ₂ LNO	Fuc ₂ Hex ₅ HexNAC ₃
3094	3116	Fuc ₃ LNO	Fuc ₃ Hex ₅ HexNAC ₃
3209	3231	Fuc ₁ LND	FucHex ₆ HexNAC ₄
Acidic peracetylated oligosaccharides			
1036	1058	NeuAcLac (Lacton)	NeuAcHex ₂
1096	1118	NeuAcLac	NeuAcHex ₂
1671	–	NeuAcLNT	NeuAcHex ₃ HexNAC
2028	–	NeuAc ₂ LNT (Lacton)	NeuAc ₂ Hex ₃ HexNAC
2246	–	NeuAcLNH	NeuAcHex ₄ HexNAC ₂
2476	–	NeuAcFuc ₁ LNH	NeuAcFucHex ₄ HexNAC ₂

Lac, lactose; Hex, hexose; NAc, *N*-acetyl; FucLac, fucosyl-lactose; LNT, lacto-*N*-tetraose; FucLNT, fucosyl-lacto-*N*-tetraose; FucLNH, fucosyl-lacto-*N*-hexaose; NeuAcLac, *N*-acetylneuraminyllactose; NeuAcLNT, *N*-acetylneuraminyllacto-*N*-tetraose.

N-acetylneuraminyllacto-*N*-tetraose, were also present. In addition, fucosyl-lactosamin and fucosyl-lacto-*N*-triose were identified, with glucose split off from the reducing end of the oligosaccharides (Table 2).

Quantification of lacto-*N*-tetraose and lacto-*N*-fucopentaose II in milk and in the infants' urine

As the milk volume ingested by an infant per suckling was known, we were able to determine in one infant the total amount of LNT and LNFP II taken up with each suckling



Fig. 2. Excretion of free urinary galactose (□) and ¹³C-enrichment (■) in fractions of an infant's urine over 24 h.

Table 2. Fast atom bombardment-MS of neutral and acidic oligosaccharides from an infant's urine

Nominal m/z values of detected molecular ions			
$[M + H]^+$	$[M + Na]^+$	Oligosaccharide	Composition
Neutral peracetylated oligosaccharides			
679	701	Lac	Hex ₂
908	–	FucLacNAC	FucHexHexNAC
909	931	FucLac	FucHex ₂
1139	–	Fuc ₂ Lac	Fuc ₂ Hex ₂
1196	1218	Fuc ₁ LNTri	FucHex ₂ HexNAC
1254	1276	LNT	Hex ₃ HexNAC
1484	1506	Fuc ₁ LNT	FucHex ₃ HexNAC
1714	1736	Fuc ₂ LNT	Fuc ₂ Hex ₃ HexNAC
	1853	Hex ₆	
2289	2311	Fuc ₂ LNH	Fuc ₂ Hex ₄ HexNAC ₂
2519	2541	Fuc ₃ LNH	Fuc ₃ Hex ₄ HexNAC ₂
Native acidic oligosaccharides			
$[M - H]^-$			
632	–	NeuAcLac	NeuAcHex ₂
997	–	NeuAcLNT	NeuAcHex ₃ HexNAC
1347	–	NeuAcFuc ₁ HexNACLNT	NeuAcFucHex ₃ HexNAC ₂

Lac, lactose; Hex, hexose; FucLacNAC, fucosyl-lactosamin; FucLac, fucosyl-lactose; FucLNTri, fucosyl-lacto-*N*-triose; LNT, lacto-*N*-tetraose; NAC, *N*-acetyl; FucLNT, fucosyl-lacto-*N*-tetraose; LNHi, lacto-*N*-hexaose; FucLNHi, fucosyl-lacto-*N*-hexaose; FucLNO, fucosyl-lacto-*N*-octaose; FucLND, fucosyl-lacto-*N*-decaose; NeuAcLac, *N*-acetylneuraminyllactose; NeuAcLNT, *N*-acetylneuraminyllacto-*N*-tetraose; NeuAcFucLNHi, *N*-acetylneuraminyll-fucosyl-lacto-*N*-hexaose.

(Table 3). The total intake of both components varies between 46 and 98 mg for LNT and between 63 and 157 mg for LNFP II per suckling. The determination of the same components in the urine fraction allowed us to quantify the total urinary excretion of these HMO (Table 4).

Discussion

There is increasing evidence, supported in recent years by animal studies, that HMO have specific beneficial effects on the infant^(14,15). Concomitantly with these observations, progress in biotechnology today allows to produce at least some of the major milk oligosaccharides to be potentially added to infant formula at reasonable costs. However, to be able to decide which compound should be added, and in which concentrations or combinations, studies are needed regarding absorption, metabolism and functions in the infants. Previously, we have shown that a major part of a ¹³C-Gal bolus orally given to lactating mothers was immediately transported to the lactating gland and is directly incorporated into HMO⁽¹⁶⁾. These results have been confirmed in the present study. Thus, using stable isotopes, an *in vivo* ¹³C-labelling of HMO can be achieved to address important questions with regard to their metabolic fate.

Here, we show that in urine of ten infants, HMO are found in all samples collected within 36 h after a ¹³C-Gal bolus was given to their mothers. The infants' urinary oligosaccharides can only be derived from milk as the biosynthesis of HMO occurs exclusively in the lactating mammary gland. Also, the excretion of intact HMO in the present study with

Table 3. Milk concentration and total intake of lacto-*N*-tetraose (LNT) and lacto-*N*-fucopentaose II (LNFP II) per suckling in one infant (Mean values and standard deviations from five samples of one day)

Suckling	Volume (ml)	Concentration (mg/ml)*				Total intake through milk (mg)	
		LNT		LNFP II		LNT	LNFP II
		Mean	SD	Mean	SD		
1	97.0	0.77	0.01	1.10	0.03	74.73	106.30
2	126.1	0.78	0.01	1.25	0.04	98.22	157.17
3	58.2	0.79	0.00	1.09	0.01	46.20	63.53
4	97.0	0.77	0.09	1.07	0.11	75.08	103.80
5	97.0	0.57	0.00	0.85	0.01	55.56	82.25

term infants at 3–6 months postpartum verifies our earlier data in preterm infants showing very similar data without using stable isotopes⁽¹⁹⁾.

Besides intact HMO, we also detected cleavage products in which not, as expected, the terminal Gal but the glucose moiety on the reducing end had been split off. At the moment, no plausible explanation can be given for such an unusual metabolic degradation step. There is only one congress contribution from Lundblad's group on the renal excretion of oligosaccharides in a few preterm and full-term infants using mass spectrometric methods for analysis⁽²⁰⁾. The authors found that small 'typical milk oligosaccharides' were excreted by human milk-fed infants. Most of the analysed urine samples contained 2'- and 3-FucLac as well as Fuc₂Lac. They could not detect these components in the urine of non-breastfed infants. A further comparison with our data regarding complex oligosaccharides such as LNT and fucosylated and/or sialylated derivatives as well as NeuAcLac is not feasible, since Lundblad and co-workers did not report the analysis of such components.

There are two possible explanations for the occurrence of 'modified' oligosaccharides in urine; either they may originate from human milk itself or they are synthesised endogenously from smaller precursors. Although there is no evidence from *in vivo* studies yet, these modifications may derive from bacterial activity within the gut. Also, if such HMO modifications occur within the colon, the underlying mechanism of the succeeding absorptive process needs to be demonstrated as well.

In the present study, we also determined the amount of HMO an exclusively breastfed infant may receive with each

Table 4. Urinary concentration and total excretion of lacto-*N*-tetraose (LNT) and lacto-*N*-fucopentaose II (LNFP II) per suckling in one infant (Mean values and standard deviations from five samples of one day)

Urine sample	Volume (ml)	Concentration (μg/ml)				Total excretion in urine (mg)	
		LNT		LNFP II		LNT	LNFP II
		Mean	SD	Mean	SD		
1	35.0	13.4	0.2	13.1	0.4	0.47	0.46
2	25.0	21.2	0.5	22.6	1.5	0.53	0.56
3	26.3	11.4	0.2	12.4	0.5	0.30	0.33
4	19.7	7.3	0.1	7.0	0.1	0.14	0.14
5	23.2	11.6	0.3	13.3	0.1	0.27	0.31

suckling. We showed that the intake of individual oligosaccharides by the infant ranges from 50 to 150 mg per suckling. This large amount of oligosaccharides which 'rinse' the whole digestive tract emphasises the potential of anti-adhesive or cellular effects shown for individual carbohydrates in many *in vitro* studies^(2,9,11,21–24). However, to exert systemic effects, HMO must be absorbed and transported in the peripheral blood to specific cells where they might also be metabolised. Although HMO are considered to be indigestible⁽²⁵⁾, we detected specific components like LNT and LNFP II in the infants' urine indicating intestinal absorption. As the renal excretion was in the range between 1 and 3 mg/d, the actual intestinal absorption must be much higher, and hence, larger amounts of these HMO must have been circulating in the infant's blood. Therefore, in addition to local functions of HMO within the gastrointestinal tract, systemic effects such as the adhesion of leucocytes to endothelial cells or the interaction of platelets with neutrophils may be influenced^(12,23). Recently, an impact of HMO on brain glycoconjugate composition has also been discussed. Carlson & House⁽²⁶⁾ compared an intra-peritoneal administration to an intra-gastrical application of NeuAc on rat brain composition, and found that both oral and systemic routes resulted in significantly more cerebral and cerebellar glycolipid and glycoprotein NeuAc than did glucose injections. Compared to free NeuAc, orally given NeuAcLac, the major acidic oligosaccharide in human milk, was even more effective on brain composition. These data supported an earlier observation by Witt *et al.*⁽²⁷⁾ comparing radiolabelled free NeuAc and NeuAcLac; the authors showed a preferential incorporation of ¹⁴C-NeuAcLac in rat brain gangliosides. Our data suggest that the absorption of HMO in breastfed infants supports the possibility that sialic acid from HMO is utilised as a substrate for the biosynthesis of components including gangliosides or glycoproteins (e.g. for the brain).

In conclusion, we have shown that the application of ¹³C-labelled HMO alleviates investigations regarding the metabolic fate of HMO in human subjects. The intake of HMO by the infants ranges within several hundred mg per suckling. Some of these components are excreted as intact oligosaccharides or as slightly modified metabolites within the infants' urine. Therefore, HMO have the potential for not only local but also systemic effects. Such effects, however, remain to be demonstrated in future studies. The enormous biotechnological progress in recent years will enable necessary studies *in vivo*.

Acknowledgements

The authors appreciate the contribution of all mothers who participated in this study which was supported by the German Research Foundation DFG (Ku 781/8-3 and Ru 529/7-3). The authors have no conflict of interest. The authors' responsibilities were as follows: S. R. designed the study, supervised the analyses and wrote the paper; G. P. performed FAB-MS; C. B. performed IR-MS; M. J. L. recruited the subjects; C. K. designed the study, recruited the subjects and wrote the paper.

References

- Thurl S, Munzert M, Henker J, *et al.* (2010) Variation of human milk oligosaccharides in relation to milk groups and lactational periods. *Br J Nutr* **104**, 1261–1271.
- Kunz C, Rudloff S, Baier W, *et al.* (2000) Oligosaccharides in human milk. Structural, functional and metabolic aspects. *Ann Rev Nutr* **20**, 699–722.
- Le Pendu J (2004) Histo-blood group antigen and human milk oligosaccharides: genetic polymorphism and risk of infectious diseases. *Adv Exp Med Biol* **554**, 135–143.
- Stahl B, Thurl S, Zeng J, *et al.* (1994) Oligosaccharides from human milk as revealed by matrix-assisted laser desorption/ionization mass spectrometry. *Anal Biochem* **223**, 218–226.
- Urashima T, Kitaoka M, Terabayashi T, *et al.* (2011) Milk oligosaccharides. In *Oligosaccharides: Sources, Properties and Applications*, pp. 1–77 [NG Gordon, editor]. New York: Nova Science Publishers.
- György P, Norris RF & Rose CS (1954) Bifidus factor I. A variant of *Lactobacillus bifidus* requiring a special growth factor. *Arch Biochem Biophys* **48**, 193–201.
- Sela DA, Chapman J, Adeuya A, *et al.* (2008) The genome sequence of *Bifidobacterium longum* subsp. *infantis* reveals adaptations for milk utilization within the infant microbiome. *Proc Natl Acad Sci U S A* **105**, 18964–18969.
- Sela DA & Mills DA (2010) Nursing our microbiota: molecular linkages between bifidobacteria and milk oligosaccharides. *Trends Microbiol* **18**, 298–307.
- Sharon N & Ofek I (2000) Safe as mother's milk: carbohydrates as future anti-adhesion drugs for bacterial diseases. *Glycoconj J* **17**, 659–664.
- Angeloni S, Ridet JL, Kusy N, *et al.* (2005) Glycoprofiling with micro-arrays of glycoconjugates and lectins. *Glycobiology* **15**, 31–41.
- Coppa GV, Zampini L, Galeazzi T, *et al.* (2006) Human milk oligosaccharides inhibit the adhesion to Caco-2 cells of diarrheal pathogens: *Escherichia coli*, *Vibrio cholerae*, and *Salmonella ftyris*. *Pediatr Res* **59**, 377–382.
- Bode L, Rudloff S, Kunz C, *et al.* (2004) Human milk oligosaccharides reduce platelet–neutrophil complex formation leading to a decrease in neutrophil $\beta 2$ integrin expression. *J Leukocyte Biol* **76**, 820–826.
- Mysore JV, Wigginton T, Simon PM, *et al.* (1999) Treatment of *Helicobacter pylori* infection in rhesus monkeys using a novel antiadhesion compound. *Gastroenterology* **117**, 1316–1325.
- Wang B, Yu B, Karim M, *et al.* (2007) Dietary sialic acid supplementation improves learning and memory in piglets. *Am J Clin Nutr* **85**, 561–569.
- Fuhrer A, Sprenger N, Kurakevich E, *et al.* (2010) Milk sialyl-lactose influences colitis in mice through selective intestinal bacterial colonization. *J Exp Med* **207**, 2843–2854.
- Rudloff S, Obermeier S, Borsch C, *et al.* (2006) Incorporation of orally applied ^{13}C -galactose into milk lactose and oligosaccharides. *Glycobiology* **16**, 477–487.
- Obermeier S, Rudloff S, Pohlentz G, *et al.* (1999) Secretion of ^{13}C -labelled oligosaccharides into human milk and infant's urine after an oral ^{13}C -galactose load. *Isotopes Environ Health Stud* **35**, 119–125.
- Kunz C, Rudloff S, Hintermann A, *et al.* (1996) High-pH anion exchange chromatography with pulsed amperometric detection and molar response factors of human milk oligosaccharides. *J Chromatogr B* **685**, 211–221.
- Rudloff S, Pohlentz G, Diekmann L, *et al.* (1996) Urinary excretion of lactose and oligosaccharides in preterm infants fed human milk or infant formula. *Acta Paediat* **85**, 598–603.
- Chester MA, Lundbland A, Renlund M, *et al.* (1981) Urinary excretion of oligosaccharides by premature and full-term babies and adults. In *Proceedings of the 6th International Symposium on Glycoconjugates*, p. 213A [T Yamakawa, T Osawa and S Handa, editors]. Tokyo: Japan Scientific Societies Press Tokyo.
- Kuntz S, Rudloff S & Kunz C (2008) Oligosaccharides from human milk influence growth-related characteristics of intestinally transformed and non-transformed intestinal cells. *Br J Nutr* **99**, 462–471.
- Kuntz S, Kunz C & Rudloff S (2009) Oligosaccharides from human milk induce growth arrest via G2/M by influencing growth-related cell cycle genes in intestinal epithelial cells. *Br J Nutr* **101**, 1306–1315.
- Bode L, Kunz C, Muhly-Reinholz M, *et al.* (2004) Inhibition of monocyte, lymphocyte, and neutrophil adhesion to endothelial cells by human milk oligosaccharides. *Thromb Haemost* **92**, 1402–1410.
- Donovan SM (2009) Human milk oligosaccharides – the plot thickens. *Br J Nutr* **101**, 1267–1269.
- Gnoth MJ, Kunz C, Kinne-Saffran E, *et al.* (2000) Human milk oligosaccharides are minimally digested *in vitro*. *J Nutr* **130**, 3014–3020.
- Carlson SE & House SG (1986) Oral and intraperitoneal administration of *N*-acetylneuraminic acid: effect on rat cerebral and cerebellar *N*-acetylneuraminic acid. *J Nutr* **116**, 881–886.
- Witt W, von Nicolai H & Zilliken F (1979) Uptake and distribution of orally applied *N*-acetyl-(^{14}C)-neuraminylactose and *N*-acetyl-(^{14}C)-neuraminic acid in the organs of newborn rats. *Nutr Metab* **23**, 51–61.